

Three-quark ground-state potential in the SU(3) lattice QCD

H. Suganuma^{a *}, H. Matsufuru,^a Y. Nemoto^a and T.T. Takahashi^a

^aRCNP, Osaka University, Mihogaoka 10-1, Ibaraki 567-0049, Japan

With the smearing technique, the three-quark (3Q) ground-state potential V_{3Q} is numerically extracted in the SU(3)_c lattice QCD Monte Carlo simulation with $12^3 \times 24$ and $\beta = 5.7$ at the quenched level. With accuracy better than a few %, V_{3Q} is well described by a sum of a constant C_{3Q} , the two-body Coulomb part $-A_{3Q} \sum_{i<j} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|}$ and the three-body linear confinement part $\sigma_{3Q} L_{\min}$, where $L_{\min}$ denotes the minimal length of the color flux tube linking the three quarks. By comparing with the Q- $\bar{Q}$ potential, we find a universal feature of the string tension as $\sigma_{3Q} \simeq \sigma_{Q\bar{Q}}$ and the one-gluon-exchange result for the Coulomb coefficient as $A_{3Q} \simeq \frac{1}{2}A_{Q\bar{Q}}$. All our results including the constant term are consistent with the requirement on the diquark limit in the lattice regularization.

1. The 3Q Wilson Loop and the 3Q Ground-State Potential in QCD

Similar to the relevant role of the Q- $\bar{Q}$ potential for meson properties, the three-quark (3Q) potential [1–3] is directly responsible to the structure and properties of baryons. In spite of the importance of the 3Q potential in the hadron physics, there were only a few preliminary lattice-QCD studies for the 3Q potential done in 80's [4,5].

The 3Q ground-state potential V_{3Q} can be measured in a gauge-invariant manner using the 3Q Wilson loop W_{3Q} as shown in Fig.1,

$$V_{3Q} = - \lim_{T \rightarrow \infty} \frac{1}{T} \ln \langle W_{3Q} \rangle, \quad W_{3Q} \equiv \frac{1}{3!} \varepsilon_{abc} \varepsilon_{a'b'c'} U_1^{aa'} U_2^{bb'} U_3^{cc'}, \quad U_k \equiv \text{Pe}^{ig \int_{\Gamma_k} dx^\mu A_\mu(x)}, \quad (1)$$

similar to the derivation of the Q- $\bar{Q}$ potential from the Wilson loop. In principle, V_{3Q} can be obtained in the large T limit, however, the practical lattice-QCD calculation of $\langle W_{3Q} \rangle$ becomes severe for large T , because $\langle W_{3Q} \rangle$ decreases exponentially with T .

Physically, the true ground state of the 3Q system is expected to consist of three flux tubes, and the 3Q state expressed by the three strings generally includes many excited-state components such as flux-tube vibrational modes. Therefore, for the accurate measurement of the 3Q ground-state potential V_{3Q} , the ground-state enhancement or the excitation-component reduction by the smearing technique [1,6,7] is practically indispensable. (This smearing was not applied in Refs. [4,5].)

In this paper, we study the 3Q ground-state potential V_{3Q} using the ground-state enhancement by the gauge-covariant smearing technique for the link-variable in the SU(3)_c lattice QCD with the standard action with $\beta=5.7$ ($a \simeq 0.19\text{fm}$) and $12^3 \times 24$ at the quenched level [1]. We consider 16 patterns of the 3Q configuration where the three

*Present address: Tokyo Institute of Technology, Ohokayama 2-12-1, Meguro, Tokyo 152-8551, Japan

quarks are put on $(i, 0, 0)$, $(0, j, 0)$ and $(0, 0, k)$ in $\mathbf{R}^3$ with $0 \leq i, j, k \leq 3$ in the lattice unit. The junction point O , which does not affect V_{3Q} , is set at the origin $(0, 0, 0)$ in $\mathbf{R}^3$.

2. The Smearing Technique and the Ground-State Enhancement

The smearing technique is actually successful for accurate measurements of the $Q\bar{Q}$ potential in the lattice QCD [7]. The standard smearing for link-variables is performed by the iterative replacement of the spatial link-variable $U_i(s)$ ($i = 1, 2, 3$) by the obscured link-variable $\bar{U}_i(s) \in \text{SU}(3)_c$ [6,7] which maximizes

$$\text{Re tr} \left\{ \bar{U}_i^\dagger(s) [\alpha U_i(s) + \sum_{j \neq i} \sum_{\pm} U_{\pm j}(s) U_i(s \pm \hat{j}) U_{\pm j}^\dagger(s + \hat{i})] \right\}, \quad U_{-\mu}(s) \equiv U_\mu^\dagger(s - \hat{\mu}), \quad (2)$$

with a real smearing parameter α . The n -th smeared link-variable $U_\mu^{(n)}(s)$ is defined as

$$U_i^{(n)}(s) \equiv \bar{U}_i^{(n-1)}(s) \quad (i = 1, 2, 3), \quad U_4^{(n)}(s) \equiv U_4(s), \quad U_\mu^{(0)}(s) \equiv U_\mu(s). \quad (3)$$

As an important feature, this smearing procedure keeps the gauge covariance of the smeared link-variable $U_\mu^{(n)}(s)$ properly. In fact, the gauge-transformation property of $U_\mu^{(n)}(s)$ is just the same as that of the original link-variable $U_\mu(s)$, and therefore the gauge invariance of $O(U_\mu^{(n)}(s))$ is ensured whenever $O(U_\mu(s))$ is gauge invariant.

Since the smeared link-variable $U_\mu^{(n)}(s)$ includes a spatial extension, the “line” expressed with $U_\mu^{(n)}(s)$ physically corresponds to a “flux tube” with a spatial extension. Therefore, if a suitable smearing is done, the “line” of the smeared link-variable is expected to be close to the ground-state flux tube. Here, the overlap between the ground-state operator and the 3Q-state operator at $t = 0, T$ in the 3Q Wilson loop W_{3Q} is estimated by

$$C_0 \equiv \langle W_{3Q}(T) \rangle^{T+1} / \langle W_{3Q}(T+1) \rangle^T \in [0, 1]. \quad (4)$$

To get the ground-state-dominant 3Q system, we investigate the ground-state overlap C_0 for the 3Q Wilson loop composed of the smeared link-variable $U_\mu^{(n)}(s)$ in the lattice QCD, and we adopt the smearing with $\alpha = 2.3$ and the iteration number $n = 12$, which largely enhances C_0 for all of the 3Q configurations in consideration as shown in Fig.2.

3. Theoretical Consideration for the 3Q Ground-State Potential

In this section, we consider the potential form of V_{3Q} based on QCD. In the short-distance limit, the perturbative QCD is applicable and the Coulomb-type potential appears as the one-gluon-exchange (OGE) result. In the long-distance limit at the quenched level, the flux-tube picture would be applicable from the argument of the strong-coupling QCD [2,3,8], and hence a linear-type confinement potential is expected to appear. Actually, the $Q\bar{Q}$ potential $V_{Q\bar{Q}}$ is well reproduced by the Coulomb-plus-linear potential in the lattice QCD. Then, we conjecture that the 3Q ground-state potential V_{3Q} is also expressed by a sum of the short-distance OGE result and the long-distance flux-tube result as

$$V_{3Q} = -A_{3Q} \sum_{i < j} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|} + \sigma_{3Q} L_{\min} + C_{3Q}, \quad V_{Q\bar{Q}}(r) = -\frac{A_{Q\bar{Q}}}{r} + \sigma_{Q\bar{Q}} r + C_{Q\bar{Q}}, \quad (5)$$

where $L_{\min}$ denotes the minimal length of the total flux tubes linking the three quarks.




Figure 1. The 3Q Wilson loop W_{3Q} . The 3Q state is generated at $t = 0$ and is annihilated at $t = T$. The three quarks are spatially fixed in $\mathbf{R}^3$ for $0 < t < T$.




Figure 2. The ground-state overlap C_0 for the 3Q system with smeared link-variables (upper data) and with unsmeared link-variables (lower data). The horizontal axis denotes $L_{\min}$. For each 3Q configuration, C_0 is largely enhanced as $0.8 < C_0 < 1$ by the smearing.

Next, we consider the diquark limit, where the 3Q system becomes equivalent to a $Q\bar{Q}$ system, which leads to a physical requirement on the relation between V_{3Q} and $V_{Q\bar{Q}}$. Here, the constant term is to be considered carefully, because there appears a divergence from the Coulomb term in V_{3Q} as $\frac{-A_{3Q}}{|\mathbf{r}_i - \mathbf{r}_j|} \rightarrow -\infty$ in the continuum diquark limit. In the lattice regularization, this ultraviolet divergence is regularized to be a finite constant with the lattice spacing a as $\frac{-A_{3Q}}{|\mathbf{r}_i - \mathbf{r}_j|} \rightarrow \frac{-A_{3Q}}{\omega a}$, where ω is a dimensionless constant satisfying $0 < \omega < 1$ and $\omega \sim 1$. Thus, we get the diquark-limit requirement on the lattice,

$$\sigma_{3Q} = \sigma_{Q\bar{Q}}, \quad A_{3Q} = \frac{1}{2}A_{Q\bar{Q}}, \quad C_{3Q} - C_{Q\bar{Q}} = \frac{A_{3Q}}{\omega a} \quad (> 0). \quad (6)$$

4. Lattice QCD Results for the 3Q Ground-State Potential

We measure the 3Q ground-state potential V_{3Q} from 210 gauge configurations in the $SU(3)_c$ lattice QCD using the smearing technique, and compare the lattice data with the theoretical form of Eq.(5). Table 1 shows the best fit coefficients in V_{3Q} and $V_{Q\bar{Q}}$ in Eq.(5). Table 2 shows the comparison between the lattice QCD data V_{3Q}^{latt} and the fitting function V_{3Q}^{fit} in Eq.(5) with the three coefficients listed in Table 1.

As shown in Table 2, the three-quark ground-state potential V_{3Q} is well described by Eq.(5) with accuracy better than a few %. From Table 1, we find a universal feature of the string tension, $\sigma_{3Q} \simeq \sigma_{Q\bar{Q}}$, as well as the OGE result for the Coulomb coefficient, $A_{3Q} \simeq \frac{1}{2}A_{Q\bar{Q}}$. All the diquark-limit requirements in Eq.(6) are satisfied for $\omega \simeq 0.46$.

Another fitting with the Δ -type flux-tube ansatz [4,5] seems rather worse, because of unacceptably large $\chi^2/N_{\text{DS}} \geq 10.9$. However, as an approximation, V_{3Q} seems described by a simple sum of an effective two-body Q - Q potential with a reduced string tension as $\sigma_{QQ} \simeq 0.53\sigma$. This reduction factor can be naturally understood as a geometrical factor rather than the color factor, since the ratio between $L_{\min}$ and the perimeter length L_P satisfies $\frac{1}{2} \leq \frac{L_{\min}}{L_P} \leq \frac{1}{\sqrt{3}}$, which leads to $L_{\min}\sigma = L_P\sigma_{QQ}$ with $\sigma_{QQ} = (0.5 \sim 0.58)\sigma$.

Table 1

The string tension, the Coulomb coefficient and the constant term for the 3Q ground-state potential V_{3Q} and the Q- $\bar{Q}$ potential $V_{Q\bar{Q}}$ in the lattice unit with $a \simeq 0.19\text{fm}$.

	σ	A	C
3Q	0.1528(27)	0.1316(62)	0.9140(201)
$Q\bar{Q}$	0.1629(47)	0.2793(116)	0.6293(161)

Table 2

The lattice QCD result for the 3Q ground-state potential V_{3Q}^{latt} and the fitting function V_{3Q}^{fit} in Eq.(5) for the 3Q system where the three quarks are put on $(i, 0, 0)$, $(0, j, 0)$ and $(0, 0, k)$ in $\mathbf{R}^3$ in the lattice unit. The error of V_{3Q}^{latt} denotes the statistical error estimated with the jackknife method. (The systematic error is not included.)

(i, j, k)	V_{3Q}^{latt}	V_{3Q}^{fit}	$V_{3Q}^{\text{latt}} - V_{3Q}^{\text{fit}}$
(0, 1, 1)	0.8459(36)	0.8529	0.0070
(0, 1, 2)	1.0970(40)	1.1023	0.0053
(0, 1, 3)	1.2935(39)	1.2926	-0.0009
(0, 2, 2)	1.3164(40)	1.3262	0.0098
(0, 2, 3)	1.5032(58)	1.5069	0.0037
(0, 3, 3)	1.6741(40)	1.6808	0.0067
(1, 1, 1)	1.0231(38)	1.0091	-0.0140
(1, 1, 2)	1.2181(61)	1.2145	-0.0036
(1, 1, 3)	1.4154(49)	1.3958	-0.0196
(1, 2, 2)	1.3870(46)	1.3887	0.0017
(1, 2, 3)	1.5588(60)	1.5580	-0.0008
(1, 3, 3)	1.7141(43)	1.7195	0.0054
(2, 2, 2)	1.5216(33)	1.5230	0.0014
(2, 2, 3)	1.6745(11)	1.6755	0.0010
(2, 3, 3)	1.8242(54)	1.8169	-0.0073
(3, 3, 3)	1.9607(92)	1.9438	-0.0169

REFERENCES

1. T.T. Takahashi, H. Matsufuru, Y. Nemoto and H. Suganuma, Proc. of the TMU-Yale Symp. on "Dynamics of Gauge Fields", TMU, Tokyo, Dec. 1999, A. Chodos et al. (eds.), Universal Academy Press, Tokyo (2000) in press.
2. S. Capstick and N. Isgur, Phys. Rev. **D34** (1986) 2809.
3. N. Brambilla, G.M. Prosperi and A. Vairo, Phys. Lett. **B362** (1995) 113.
4. R. Sommer and J. Wosiek, Phys.Lett.**149B** (1984) 497; Nucl.Phys.**B267** (1986) 531.
5. H.B. Thacker, E. Eichten and J.C. Sexton, Nucl. Phys. **B** (Proc. Suppl.) **4** (1988) 234.
6. APE Collaboration, M. Albanese et al., Phys. Lett. **B192** (1987) 163.
7. G.S. Bali, C. Schlichter and K. Schilling, Phys. Rev. **D51** (1995) 5165.
8. J. Kogut and L. Susskind, Phys. Rev. **D11** (1975) 395.